

The University of the State of New York
REGENTS HIGH SCHOOL EXAMINATION

GEOMETRY

Friday, June 21, 2019 — 9:15 a.m. to 12:15 p.m.


MODEL RESPONSE SET

Table of Contents

Question 25	2
Question 26	7
Question 27	11
Question 28	16
Question 29	21
Question 30	28
Question 31	33
Question 32	39
Question 33	55
Question 34	63
Question 35	71

Question 25

25 Triangle $A'B'C'$ is the image of triangle ABC after a dilation with a scale factor of $\frac{1}{2}$ and centered at point A . Is triangle ABC congruent to triangle $A'B'C'$? Explain your answer.



No $\triangle ABC$ isn't congruent to $\triangle A'B'C'$ because a dilation doesn't preserve segment lengths

Score 2: The student gave a complete and correct response.

Question 25

25 Triangle $A'B'C'$ is the image of triangle ABC after a dilation with a scale factor of $\frac{1}{2}$ and centered at point A . Is triangle ABC congruent to triangle $A'B'C'$? Explain your answer.

no, they're not
congruent b/c
 $A'B'C'$ had a
dilation of $\frac{1}{2}$
so the image will
be a lot smaller
than ABC . It will be
the same shape but
different size.

Score 2: The student gave a complete and correct response.

Question 25

25 Triangle $A'B'C'$ is the image of triangle ABC after a dilation with a scale factor of $\frac{1}{2}$ and centered at point A . Is triangle ABC congruent to triangle $A'B'C'$? Explain your answer.

No, dilations always produce similar triangles.

Score 1: The student wrote an incomplete explanation.

Question 25


25 Triangle $A'B'C'$ is the image of triangle ABC after a dilation with a scale factor of $\frac{1}{2}$ and centered at point A . Is triangle ABC congruent to triangle $A'B'C'$? Explain your answer.

No, a dilation is not a rigid motion.

Score 1: The student wrote an incomplete explanation.

Question 25

25 Triangle $A'B'C'$ is the image of triangle ABC after a dilation with a scale factor of $\frac{1}{2}$ and centered at point A . Is triangle ABC congruent to triangle $A'B'C'$? Explain your answer.



Yes $\triangle ABC$ is \cong to $\triangle A'B'C'$
by AAA.

Score 0: The student gave a completely incorrect response.

Question 26

26 Determine and state the area of triangle PQR , whose vertices have coordinates $P(-2, -5)$, $Q(3, 5)$, and $R(6, 1)$.


[The use of the set of axes below is optional.]

$$QR = \sqrt{3^2 + (-4)^2} = \sqrt{25} = 5$$

$$PR = \sqrt{8^2 + 6^2} = \sqrt{100} = 10$$

$$\begin{aligned} m_{\overline{QR}} &= -\frac{4}{3} \\ m_{\overline{PR}} &= \frac{6}{8} = \frac{3}{4} \end{aligned} \quad \left. \vphantom{\begin{aligned} m_{\overline{QR}} &= -\frac{4}{3} \\ m_{\overline{PR}} &= \frac{6}{8} = \frac{3}{4} \end{aligned}} \right\} \perp$$

$$A = \frac{1}{2}(5)(10) = 25$$



Score 2: The student gave a complete and correct response.

Question 26

26 Determine and state the area of triangle PQR , whose vertices have coordinates $P(-2, -5)$, $Q(3, 5)$, and $R(6, 1)$.

[The use of the set of axes below is optional.]

$$\text{Total} = 10 \cdot 8 = 80$$

$$B = \frac{1}{2}(3 \cdot 4)$$

$$= 6$$

$$C = \frac{1}{2}(6 \cdot 8)$$


$$= 24$$

$$D = \frac{1}{2}(5 \cdot 10)$$

$$= 25$$

$$A = 80 - 55$$

$$\Delta PQR = 25$$




Score 2: The student gave a complete and correct response.

Question 26

26 Determine and state the area of triangle PQR , whose vertices have coordinates $P(-2, -5)$, $Q(3, 5)$, and $R(6, 1)$.

[The use of the set of axes below is optional.]



Score 1: The student did not show work to determine the lengths of \overline{QR} and \overline{PR} .

Question 26

26 Determine and state the area of triangle PQR , whose vertices have coordinates $P(-2, -5)$, $Q(3, 5)$, and $R(6, 1)$.

[The use of the set of axes below is optional.]


$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$PQ = \sqrt{(-2 - 3)^2 + (5 - 5)^2} = 5$$

$$PR = \sqrt{(6 - 6)^2 + (5 - 1)^2} = 4$$

$$QR = \sqrt{(3 - 6)^2 + (5 - 1)^2} = 5$$


$$A = 1450$$



Score 0: The student did not show enough correct relevant work to receive any credit.

Question 27

27 A support wire reaches from the top of a pole to a clamp on the ground. The pole is perpendicular to the level ground and the clamp is 10 feet from the base of the pole. The support wire makes a 68° angle with the ground. Find the length of the support wire to the nearest foot.



$$\cos = \frac{A}{H}$$

$$\frac{\cos(68)}{1} = \frac{10}{x}$$

~~$$\cos(68)x = 10$$~~

$$\frac{\cos(68)x}{\cos(68)} = \frac{10}{\cos(68)}$$

$$x = 26.69467163$$


support wire = 27 ft.

Score 2: The student gave a complete and correct response.

Question 27

27 A support wire reaches from the top of a pole to a clamp on the ground. The pole is perpendicular to the level ground and the clamp is 10 feet from the base of the pole. The support wire makes a 68° angle with the ground. Find the length of the support wire to the nearest foot.

SIN CAT C



$$\frac{\tan 68^\circ}{1} = \frac{x}{10}$$

$$x = 24.75$$

$$10^2 + 24.75^2 = c^2$$

$$100 + 612.5625 = c^2$$


$$712.5625 = c^2$$

$$c = 27$$

Score 2: The student gave a complete and correct response.

Question 27

27 A support wire reaches from the top of a pole to a clamp on the ground. The pole is perpendicular to the level ground and the clamp is 10 feet from the base of the pole. The support wire makes a 68° angle with the ground. Find the length of the support wire to the nearest foot.



$$\cos 68^\circ = \frac{10}{x}$$

$$\cos 68 \cdot 10 = x$$


$$4 \approx x$$

4 feet

Score 1: The student wrote a correct trigonometric equation to find the length of the support wire, but no further correct work was shown.

Question 27

27 A support wire reaches from the top of a pole to a clamp on the ground. The pole is perpendicular to the level ground and the clamp is 10 feet from the base of the pole. The support wire makes a 68° angle with the ground. Find the length of the support wire to the nearest foot.



$$\frac{\sin(68)}{1} = \frac{10}{X}$$

$$X \sin(68) = 10$$
$$\frac{X \sin(68)}{\sin(68)} = \frac{10}{\sin(68)}$$


$$X \approx 11 \text{ ft}$$

the wire to the clamp is approximately 11 ft.

Score 1: The student showed appropriate work based on an incorrect location of the angle of elevation.

Question 27

27 A support wire reaches from the top of a pole to a clamp on the ground. The pole is perpendicular to the level ground and the clamp is 10 feet from the base of the pole. The support wire makes a 68° angle with the ground. Find the length of the support wire to the *nearest foot*.



$$\sin 68 = \frac{10}{x}$$


$$x = \frac{10}{\sin 68}$$

$$x = 10.8 \text{ feet}$$

Score 0: The student labeled the diagram incorrectly and did not round the answer to the nearest foot.

Question 28

28 In the diagram below, circle O has a radius of 10.



If $m\widehat{AB} = 72^\circ$, find the area of shaded sector AOB , in terms of π .

$$\frac{72}{360} = \frac{x}{100\pi}$$


$$7200\pi = 360x$$

$$20\pi = x$$

Score 2: The student gave a complete and correct response.

Question 28

28 In the diagram below, circle O has a radius of 10.



If $m\widehat{AB} = 72^\circ$, find the area of shaded sector AOB , in terms of π .

$$\frac{72}{360} = \frac{1}{5}$$

$$\begin{aligned} \text{Area circle} &= \pi r^2 \\ &= \pi 10^2 \end{aligned}$$


$$A = 100\pi$$

$$\frac{1}{5} (100\pi) = \boxed{20\pi}$$

Score 2: The student gave a complete and correct response.

Question 28

28 In the diagram below, circle O has a radius of 10.



If $m\widehat{AB} = 72^\circ$, find the area of shaded sector AOB , in terms of π .

$$A_{\text{sector}} = \frac{\cancel{4}}{360} \pi r^2$$


$$A_{\text{sector}} = \frac{72}{360} \pi 100$$

$$A_{\text{sector}} = 62.8$$

Score 1: The student wrote an appropriate answer, but not in terms of π .

Question 28

28 In the diagram below, circle O has a radius of 10.



If $m\widehat{AB} = 72^\circ$, find the area of shaded sector AOB , in terms of π .

$$C = 2\pi r$$


$$C = 2\pi(10)$$

$$C = 20\pi$$

Score 0: The student gave a correct response that was obtained by an incorrect procedure.

Question 28

28 In the diagram below, circle O has a radius of 10.



If $m\widehat{AB} = 72^\circ$, find the area of shaded sector AOB , in terms of π .

$$A = \pi r^2$$


$$A = \pi 10^2$$

$$A = \pi 100$$

Score 0: The student did not show enough relevant work to receive any credit.

Question 29

29 On the set of axes below, $\triangle ABC \cong \triangle STU$.




Describe a sequence of rigid motions that maps $\triangle ABC$ onto $\triangle STU$.

A rotation of 90° counter clockwise about the origin.

Score 2: The student gave a complete and correct response.

Question 29

29 On the set of axes below, $\triangle ABC \cong \triangle STU$.




Describe a sequence of rigid motions that maps $\triangle ABC$ onto $\triangle STU$.

To map $\triangle ABC$ onto $\triangle STU$ first it needs to rotate 90° counterclockwise from $(-4, 2)$, next translate 6 units down and then 2 units right.

Score 2: The student gave a complete and correct response.

Question 29

29 On the set of axes below, $\triangle ABC \cong \triangle STU$.




Describe a sequence of rigid motions that maps $\triangle ABC$ onto $\triangle STU$.

Reflect $\triangle ABC$ over the y -axis, then reflect the image over the x -axis, then rotate around the origin 90° clockwise onto $\triangle STU$.

Score 2: The student gave a complete and correct response.

Question 29

29 On the set of axes below, $\triangle ABC \cong \triangle STU$.




Describe a sequence of rigid motions that maps $\triangle ABC$ onto $\triangle STU$.

A counter-clockwise rotation 90°

Score 1: The student did not state the center of rotation.

Question 29

29 On the set of axes below, $\triangle ABC \cong \triangle STU$.




Describe a sequence of rigid motions that maps $\triangle ABC$ onto $\triangle STU$.

- 1.) Rotate $\triangle STU$ 90° clockwise around point T
- 2.) Translate $\triangle S'T'U'$ along vector $\vec{U'C}$ until it maps on to $\triangle ABC$

Score 1: The student mapped $\triangle STU$ onto $\triangle ABC$.

Question 29

29 On the set of axes below, $\triangle ABC \cong \triangle STU$.




Describe a sequence of rigid motions that maps $\triangle ABC$ onto $\triangle STU$.

COUNTER CLOCKWISE ROTATION

Score 0: The student did not state the number of degrees in the rotation and did not state the center of rotation.

Question 29

29 On the set of axes below, $\triangle ABC \cong \triangle STU$.




Describe a sequence of rigid motions that maps $\triangle ABC$ onto $\triangle STU$.

$\triangle ABC$ rotated clockwise on point C and did a translation of $(2, -6)$.

Score 0: The student did not state the number of degrees in the rotation and wrote an incorrect translation from point C .

Question 30

30 In right triangle PRT , $m\angle P = 90^\circ$, altitude \overline{PQ} is drawn to hypotenuse \overline{RT} , $RT = 17$, and $PR = 15$.



Determine and state, to the *nearest tenth*, the length of \overline{RQ} .

$$\frac{h}{LL} = \frac{h}{LL}$$

$$\overline{RQ} = 13.2$$

$$\frac{17}{15} = \frac{15}{x}$$


$$\frac{17x = 225}{17}$$

$$x = 13.235 \approx 13.2$$

Score 2: The student gave a complete and correct response.


Question 30

30 In right triangle PRT , $m\angle P = 90^\circ$, altitude \overline{PQ} is drawn to hypotenuse \overline{RT} , $RT = 17$, and $PR = 15$.



Determine and state, to the *nearest tenth*, the length of \overline{RQ} .

$$\frac{8}{17} = \frac{x}{8}$$



$$\frac{17x}{17} = \frac{64}{17}$$

$$x = 3.76471$$


$$17 - 3.76471$$

$$\overline{RQ} = 13.2 \text{ units}$$

Score 2: The student gave a complete and correct response.

Question 30

30 In right triangle PRT , $m\angle P = 90^\circ$, altitude \overline{PQ} is drawn to hypotenuse \overline{RT} , $RT = 17$, and $PR = 15$.



Determine and state, to the *nearest tenth*, the length of \overline{RQ} .

$$\cos X = \frac{15}{17}$$

$$X = \cos^{-1}\left(\frac{15}{17}\right)$$

$$X = 28.07248694$$

$$\cos(28.07248694) = \frac{y}{15}$$

$$y = 15 \cdot \cos(28.07248694)$$


$$y = 13.23529412$$

13.2

Score 2: The student gave a complete and correct response.

Question 30

30 In right triangle PRT , $m\angle P = 90^\circ$, altitude \overline{PQ} is drawn to hypotenuse \overline{RT} , $RT = 17$, and $PR = 15$.



Determine and state, to the *nearest tenth*, the length of \overline{RQ} .

$$\frac{17}{15} = \frac{15}{x}$$


$$\frac{15x}{15} = \frac{255}{15}$$

$$x = 17$$

Score 1: The student wrote a correct proportion to solve for RQ , but no further correct work was shown.

Question 30

30 In right triangle PRT , $m\angle P = 90^\circ$, altitude \overline{PQ} is drawn to hypotenuse \overline{RT} , $RT = 17$, and $PR = 15$.



Determine and state, to the *nearest tenth*, the length of \overline{RQ} .

{8, 15, 17} triple

HLLS

$$\frac{17}{8} = \frac{15}{x}$$

$$\frac{17x}{17} = \frac{120}{17}$$

$$x = 7.0588$$


$$17 - 7.0588$$

$$9.9412 = QR$$

Score 0: The student wrote an incorrect proportion to solve for TQ , then made a rounding error.

Question 31


31 Given circle O with radius \overline{OA} , use a compass and straightedge to construct an equilateral triangle inscribed in circle O . [Leave all construction marks.]



Score 2: The student gave a complete and correct response.

Question 31


31 Given circle O with radius \overline{OA} , use a compass and straightedge to construct an equilateral triangle inscribed in circle O . [Leave all construction marks.]



Score 2: The student gave a complete and correct response.

Question 31


31 Given circle O with radius \overline{OA} , use a compass and straightedge to construct an equilateral triangle inscribed in circle O . [Leave all construction marks.]



Score 2: The student gave a complete and correct response.

Question 31


31 Given circle O with radius \overline{OA} , use a compass and straightedge to construct an equilateral triangle inscribed in circle O . [Leave all construction marks.]



Score 1: The student constructed an equilateral triangle, but it was not inscribed in circle O .

Question 31


31 Given circle O with radius \overline{OA} , use a compass and straightedge to construct an equilateral triangle inscribed in circle O . [Leave all construction marks.]



Score 1: The student constructed an inscribed isosceles right triangle.

Question 31

31 Given circle O with radius \overline{OA} , use a compass and straightedge to construct an equilateral triangle inscribed in circle O . [Leave all construction marks.]



Score 0: The student gave a drawing that is not a construction.

Question 32

32 Riley plotted $A(-1,6)$, $B(3,8)$, $C(6,-1)$, and $D(1,0)$ to form a quadrilateral.

Prove that Riley's quadrilateral $ABCD$ is a trapezoid.

[The use of the set of axes on the next page is optional.]

$$\begin{aligned} m_{AD} &= \frac{0-6}{1-(-1)} = \frac{-6}{2} = \frac{-3}{1} \\ m_{BC} &= \frac{-1-8}{6-3} = \frac{-9}{3} = \frac{-3}{1} \end{aligned} \quad \left. \vphantom{\begin{aligned} m_{AD} \\ m_{BC} \end{aligned}} \right\} \overline{AD} \parallel \overline{BC}$$

Riley's quadrilateral $ABCD$ is a trapezoid because it has a pair of parallel sides.

Score 4: The student gave a complete and correct response.

Question 32 continued

Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that $ABCD$ is *not* an isosceles trapezoid.


$$AC = \sqrt{(-1-6)^2 + (6--1)^2} \approx 9.899$$

$$BD = \sqrt{(8-0)^2 + (3-1)^2} \approx 8.246$$

-1, 6
3, 8

6, -1
1, 0

$ABCD$ is not an isosceles trapezoid because its diagonals aren't congruent.
 $9.899 \neq 8.246$



Question 32

32 Riley plotted $A(-1,6)$, $B(3,8)$, $C(6,-1)$, and $D(1,0)$ to form a quadrilateral.

Prove that Riley's quadrilateral $ABCD$ is a trapezoid.

[The use of the set of axes on the next page is optional.]

$$\begin{array}{l} m_{\overline{AB}} = \frac{2}{4} \\ m_{\overline{BC}} = \frac{9}{-3} = \frac{-3}{1} \\ m_{\overline{CD}} = \frac{-1}{5} \\ m_{\overline{AD}} = \frac{-6}{0} = \frac{-3}{1} \end{array}$$

Not same slope \therefore Not // sides

Same slope \therefore // sides

Since there is only 1 pair of opp sides //,
Quad $ABCD$ is a trapezoid.

Score 4: The student gave a complete and correct response.

Question 32 continued


Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that $ABCD$ is *not* an isosceles trapezoid.

$$d \overline{DB} = \sqrt{68}$$

$$d \overline{AC} = \sqrt{98}$$

diagonals have
different dist.
 \therefore They are not \cong .

Trapez $ABCD$ is Not Isosceles



Question 32

32 Riley plotted $A(-1,6)$, $B(3,8)$, $C(6,-1)$, and $D(1,0)$ to form a quadrilateral.

Prove that Riley's quadrilateral $ABCD$ is a trapezoid.

[The use of the set of axes on the next page is optional.]

Plan: Show that $ABCD$ is a trapezoid by proving that its bases are \parallel

$$\text{Slope of } \overline{AD} = \frac{6-0}{-1-1} = \frac{6}{-2} = -3$$

$$\text{Slope of } \overline{BC} = \frac{8-(-1)}{3-6} = \frac{9}{-3} = -3$$

$$-3 = -3$$

$\therefore \overline{AD} \parallel \overline{BC}$ b/c equal slopes yield \parallel lines and $ABCD$ is a trapezoid b/c it is a quadrilateral w/ at least one pair of opp. sides \parallel .

Score 3: The student made a computational error in determining the length of diagonal \overline{BD} .

Question 32 continued

Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that $ABCD$ is *not* an isosceles trapezoid.

Plan: Prove that $ABCD$ is not an isosceles trapezoid by showing that the diagonals are not \cong

$$AC = \sqrt{(-1-6)^2 + (6--1)^2}$$

$$AC = \sqrt{49 + 49}$$

$$AC = \sqrt{98}$$


$$BD = \sqrt{(3-1)^2 + (8-0)^2}$$

$$= \sqrt{4 + 64}$$

$$\Rightarrow \sqrt{70}$$

$$\sqrt{98} \neq \sqrt{70}$$

$\therefore ABCD$ is not isosceles b/c its diagonals are not \cong



Question 32

32 Riley plotted $A(-1,6)$, $B(3,8)$, $C(6,-1)$, and $D(1,0)$ to form a quadrilateral.

Prove that Riley's quadrilateral $ABCD$ is a trapezoid.

[The use of the set of axes on the next page is optional.]

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m_{\overline{BC}} = \frac{-1 - 8}{6 - 3} = \frac{-9}{3} = \boxed{-3}$$

$$m_{\overline{AD}} = \frac{0 - 6}{1 - (-1)} = \frac{-6}{2} = \boxed{-3}$$

$$m_{\overline{BC}} = m_{\overline{AD}}$$

$$\therefore \boxed{\overline{BC} \parallel \overline{AD}}$$

$ABCD$ is a trapezoid because it has one pair of parallel sides.

Score 2: The student correctly proved quadrilateral $ABCD$ is a trapezoid, but no further correct work is shown.

Question 32 continued


Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that $ABCD$ is *not* an isosceles trapezoid.

$$D = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$D_{AB} = \sqrt{(3 - (-1))^2 + (8 - 6)^2}$$

$$D_{AB} = \sqrt{4^2 + 2^2} = \sqrt{16 + 4} = \boxed{20}$$

$$D_{CD} = \sqrt{\quad}$$



Question 32

32 Riley plotted $A(-1,6)$, $B(3,8)$, $C(6,-1)$, and $D(1,0)$ to form a quadrilateral.

Prove that Riley's quadrilateral $ABCD$ is a trapezoid.

[The use of the set of axes on the next page is optional.]

$\overline{AD} \cong \overline{BC} \rightarrow$ if lines are \parallel , then
sides are \cong

Score 2: The student correctly proved the diagonals are not congruent, so quadrilateral $ABCD$ is not an isosceles trapezoid.


Question 32 continued

Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that $ABCD$ is *not* an isosceles trapezoid.

$$\begin{aligned} AC &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\ &= \sqrt{(6+1)^2 + (-1-6)^2} \\ &= \sqrt{49+49} \\ &= \sqrt{98} = 9.8994\dots \end{aligned}$$

$$\begin{aligned} BD &= \sqrt{(3-1)^2 + (8-0)^2} \\ &= \sqrt{4+64} \\ &= \sqrt{68} = 8.2462\dots \end{aligned}$$

Not isosceles due to distances not being equal



Question 32

32 Riley plotted $A(-1,6)$, $B(3,8)$, $C(6,-1)$, and $D(1,0)$ to form a quadrilateral.

Prove that Riley's quadrilateral $ABCD$ is a trapezoid.

[The use of the set of axes on the next page is optional.]

$$m = \frac{\text{rise}}{\text{run}}$$

$$m_{\overline{AD}} = \frac{-6}{2} = -\frac{3}{1}$$

$$m_{\overline{BC}} = \frac{-9}{3} = -\frac{3}{1}$$

$$m_{\overline{AD}} = m_{\overline{BC}}$$

Score 2: Appropriate work is shown, but both concluding statements are missing.


Question 32 continued

Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that $ABCD$ is *not* an isosceles trapezoid.

$$d_{\overline{AC}} = \sqrt{7^2 + 7^2} = \sqrt{98}$$

$$d_{\overline{AC}} \neq d_{\overline{BD}}$$

$$d_{\overline{BD}} = \sqrt{8^2 + 2^2} = \sqrt{68}$$



Question 32

32 Riley plotted $A(-1,6)$, $B(3,8)$, $C(6,-1)$, and $D(1,0)$ to form a quadrilateral.

Prove that Riley's quadrilateral $ABCD$ is a trapezoid.

[The use of the set of axes on the next page is optional.]

$$AD = \frac{0-6}{1+1} = \frac{-6}{2} = -3$$

Slope of AD is -3

Slope of BC is also -3


$$BC = \frac{-1-8}{6-3} = \frac{-9}{3} = -3$$

has one set of parallel lines

Score 1: The student found the slopes of bases \overline{AD} and \overline{BC} , but the concluding statement is incomplete. No further correct work is shown.

Question 32 continued

Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that $ABCD$ is *not* an isosceles trapezoid.



Question 32

32 Riley plotted $A(-1,6)$, $B(3,8)$, $C(6,-1)$, and $D(1,0)$ to form a quadrilateral.

Prove that Riley's quadrilateral $ABCD$ is a trapezoid.

[The use of the set of axes on the next page is optional.]

Quadrilateral $ABCD$ is a trapezoid


4 sides

One set of parallel lines

Score 0: The student did not show enough correct relevant work to receive any credit.

Question 32 continued


Riley defines an isosceles trapezoid as a trapezoid with congruent diagonals. Use Riley's definition to prove that $ABCD$ is *not* an isosceles trapezoid.



Both diagonals have different slopes so they are not congruent, therefore it cannot be an isosceles trapezoid.

Question 33

33 A child-sized swimming pool can be modeled by a cylinder. The pool has a diameter of $6\frac{1}{2}$ feet and a height of 12 inches. The pool is filled with water to $\frac{2}{3}$ of its height. Determine and state the volume of the water in the pool, to the nearest cubic foot.



$$V = \pi 3.25^2 \times \frac{2}{3}$$

$$\text{Volume of water} = 22.12 \\ \approx 22$$

$$\text{height: } 12''$$

$$\text{depth: } \frac{2}{3} \text{ of a foot}$$

One cubic foot equals 7.48 gallons of water. Determine and state, to the nearest gallon, the number of gallons of water in the pool.

$$7.48 \times 22.12$$

$$= 165.45$$

$$\approx 165 \text{ gallons}$$

Score 4: The student gave a complete and correct response.

Question 33

33 A child-sized swimming pool can be modeled by a cylinder. The pool has a diameter of $6\frac{1}{2}$ feet and a height of 12 inches. The pool is filled with water to $\frac{2}{3}$ of its height. Determine and state the volume of the water in the pool, to the *nearest cubic foot*.

$$\begin{aligned} V &= Bh \\ V &= (\pi \cdot 3\frac{1}{4}) \cdot 1 \\ V &= 33.18 \text{ ft}^3 \times \frac{2}{3} = 22 \text{ ft}^3 \end{aligned}$$


One cubic foot equals 7.48 gallons of water. Determine and state, to the *nearest gallon*, the number of gallons of water in the pool.

$$22 \cdot 7.48 = 165 \text{ gal}$$

Score 4: The student gave a complete and correct response.

Question 33

33 A child-sized swimming pool can be modeled by a cylinder. The pool has a diameter of $6\frac{1}{2}$ feet and a height of 12 inches. The pool is filled with water to $\frac{2}{3}$ of its height. Determine and state the volume of the water in the pool, to the *nearest cubic foot*.



$$\frac{38226.89941}{12} =$$

$$3185.574951$$

$$\textcircled{3186 \text{ ft}^3}$$

$$V = \pi r^2 h$$

$$V = \pi (39)^2 \cdot 12$$

$$V = \pi (1521) \cdot 12$$

$$V = 18252 \pi$$

$$V = \frac{57340.34911}{3}$$

$$19113.4497 \cdot 2 =$$

$$38226.89941 \text{ in}^3$$

One cubic foot equals 7.48 gallons of water. Determine and state, to the *nearest gallon*, the number of gallons of water in the pool.

$$3186 * 7.48 = 23831.28$$

$$\textcircled{2383 \text{ gallons}}$$

Score 3: The student made an error in converting cubic inches into cubic feet.

Question 33


33 A child-sized swimming pool can be modeled by a cylinder. The pool has a diameter of $6\frac{1}{2}$ feet and a height of 12 inches. The pool is filled with water to $\frac{2}{3}$ of its height. Determine and state the volume of the water in the pool, to the *nearest cubic foot*.

$$\begin{array}{l} \text{Diameter} = \text{Radius} \\ \frac{6.5}{2} = 3.25 \end{array}$$

$$\text{Volume of cylinder} = \pi r^2 h$$

$$\begin{aligned} \text{Volume} &= \pi (3.5)^2 \cdot \left(\frac{2}{3}\right) \\ &= 26 \text{ feet}^3 \end{aligned}$$

The volume of water is 26 feet³



One cubic foot equals 7.48 gallons of water. Determine and state, to the *nearest gallon*, the number of gallons of water in the pool.

$$\frac{7.48}{1} = \frac{x}{26}$$

$$\begin{array}{l} 1 = 7.48 \\ x = \end{array}$$


$$x = 194.48 \text{ gallons}$$

The number of gallons of water in this pool is 194 gallons.

Score 3: The student made a transcription error by using a radius of 3.5 instead of 3.25.

Question 33

33 A child-sized swimming pool can be modeled by a cylinder. The pool has a diameter of $6\frac{1}{2}$ feet and a height of 12 inches. The pool is filled with water to $\frac{2}{3}$ of its height. Determine and state the volume of the water in the pool, to the *nearest cubic foot*.



$$\begin{aligned}d &= 6.5 \\r &= 3.25 \\h &= \frac{2}{3}\end{aligned}$$

$$\begin{aligned}V_{\text{cyl}} &= Bh \\V_{\text{cyl}} &= \pi r^2 h\end{aligned}$$

$$\begin{aligned}V_{\text{cyl}} &= (3.25)^2 \pi \cdot \frac{2}{3} \\V &= 22.12204827 \\V &= 22.1 \text{ ft}^3\end{aligned}$$

One cubic foot equals 7.48 gallons of water. Determine and state, to the *nearest gallon*, the number of gallons of water in the pool.

$$\frac{1 \text{ ft}^3}{7.48 \text{ gal}} = \frac{22.1 \text{ ft}^3}{x}$$


$$x \text{ ft}^3 = 165.308 \text{ ft}^3 \text{ gal}$$

$$x = 165.3 \text{ gallons}$$

Score 3: The student made the same rounding error for both answers.

Question 33

33 A child-sized swimming pool can be modeled by a cylinder. The pool has a diameter of $6\frac{1}{2}$ feet and a height of 12 inches. The pool is filled with water to $\frac{2}{3}$ of its height. Determine and state the volume of the water in the pool, to the *nearest cubic foot*.



One cubic foot equals 7.48 gallons of water. Determine and state, to the *nearest gallon*, the number of gallons of water in the pool.


$22.12 / 7.48$
 2.957219251
so approx. 3 gallons

Score 2: The student made a rounding error in determining the volume and a computational error in determining the number of gallons.

Question 33

33 A child-sized swimming pool can be modeled by a cylinder. The pool has a diameter of $6\frac{1}{2}$ feet and a height of 12 inches. The pool is filled with water to $\frac{2}{3}$ of its height. Determine and state the volume of the water in the pool, to the *nearest cubic foot*.

$$\begin{aligned} V &= \pi r^2 h \\ V &= \pi 3^2 (12) \\ V &= \pi 9 (12) \\ V &= \pi 108 \\ \boxed{V &= 339 \text{ ft}^3} \end{aligned}$$



One cubic foot equals 7.48 gallons of water. Determine and state, to the *nearest gallon*, the number of gallons of water in the pool.

$$1 \text{ ft}^3 = 7.48$$

$$339 \times 7.48 = 2535.72$$

$$\boxed{2535.72 \text{ gallons of water}}$$

Score 1: The student found the volume incorrectly and made a rounding error.

Question 33

33 A child-sized swimming pool can be modeled by a cylinder. The pool has a diameter of $6\frac{1}{2}$ feet and a height of 12 inches. The pool is filled with water to $\frac{2}{3}$ of its height. Determine and state the volume of the water in the pool, to the *nearest cubic foot*.

$$\begin{aligned} V &= (6\frac{1}{2})(12)(\frac{2}{3})\pi \\ &= 52\pi \text{ ft}^3 \end{aligned}$$


One cubic foot equals 7.48 gallons of water. Determine and state, to the *nearest gallon*, the number of gallons of water in the pool.

$$\frac{52\pi}{7.48} = 21.84 \text{ gallons}$$

Score 0: The student gave a completely incorrect response.

Question 34

34 Nick wanted to determine the length of one blade of the windmill pictured below. He stood at a point on the ground 440 feet from the windmill's base. Using surveyor's tools, Nick measured the angle between the ground and the highest point reached by the top blade and found it was 38.8° . He also measured the angle between the ground and the lowest point of the top blade, and found it was 30° .



Determine and state a blade's length, x , to the nearest foot.

$$440 \cdot \tan(30) = \frac{N}{440} \cdot 440 \quad N = 254.0311184$$

$$\tan(38.8) = \frac{Y}{440} \cdot Y = 353.7690827$$


$$\begin{array}{r} 353.7690827 \\ - 254.0311184 \\ \hline 99.7379643 \end{array}$$

one blade's length is 100 ft

Score 4: The student gave a complete and correct response.

Question 34

34 Nick wanted to determine the length of one blade of the windmill pictured below. He stood at a point on the ground 440 feet from the windmill's base. Using surveyor's tools, Nick measured the angle between the ground and the highest point reached by the top blade and found it was 38.8° . He also measured the angle between the ground and the lowest point of the top blade, and found it was 30° .



Determine and state a blade's length, x , to the nearest foot.

~~100 ft~~ 100 ft.

$$\tan 30 = \frac{z}{440} \quad 440 \tan 30 = z \quad z = 254$$

$$\tan 38.8 = \frac{y}{440}$$


$$440 \tan 38.8 = y \quad y = 353.8$$

353.8
-254.0
99.8 (100)

Score 4: The student gave a complete and correct response.

Question 34

34 Nick wanted to determine the length of one blade of the windmill pictured below. He stood at a point on the ground 440 feet from the windmill's base. Using surveyor's tools, Nick measured the angle between the ground and the highest point reached by the top blade and found it was 38.8° . He also measured the angle between the ground and the lowest point of the top blade, and found it was 30° .



Determine and state a blade's length, x , to the nearest foot.

$$\begin{aligned} \tan(30) &= \frac{Y}{440} \\ Y &= 254 \end{aligned}$$

$$\begin{aligned} \cos(30) &= \frac{440}{x} \\ 508 &= x \cdot \frac{\cos(30)}{\cos(30)} \end{aligned}$$

$$\begin{aligned} \tan(38.8) &= \frac{0}{440} \\ 0 &= 440 \cdot \tan(38) \\ &= 343.7 \\ &= 344 \end{aligned}$$


$$\begin{array}{r} 344 \\ - 254 \\ \hline 90 \end{array}$$

90 ft


Score 3: The student made a transcription error by using 38° instead of 38.8° .

Question 34

34 Nick wanted to determine the length of one blade of the windmill pictured below. He stood at a point on the ground 440 feet from the windmill's base. Using surveyor's tools, Nick measured the angle between the ground and the highest point reached by the top blade and found it was 38.8° . He also measured the angle between the ground and the lowest point of the top blade, and found it was 30° .




Determine and state a blade's length, x , to the nearest foot.



$$\tan 30 = \frac{y}{440}$$

$$y = \tan 30(440)$$

$$y = -2818.3457$$



$$\tan 38.8 = \frac{x+y}{440}$$

$$\tan 38.8 = \frac{x + -2818.3457}{440}$$

$$x + -2818.3457 = \tan 38.8(440)$$

$$x + -2818.3457 = 866.3969$$


$$+ 2818.3457 \quad + 2818.3457$$

$$\boxed{x = 3685\text{ ft}}$$

Score 3: The student made an error using radian measure.

Question 34

34 Nick wanted to determine the length of one blade of the windmill pictured below. He stood at a point on the ground 440 feet from the windmill's base. Using surveyor's tools, Nick measured the angle between the ground and the highest point reached by the top blade and found it was 38.8° . He also measured the angle between the ground and the lowest point of the top blade, and found it was 30° .



Determine and state a blade's length, x , to the nearest foot.

$$\tan \angle = O/A$$

$$\tan(38.8) = y/440$$


$$y = 353.769$$

$$y = 354$$

Score 2: The student correctly found the height to the top of the top blade.


Question 34

34 Nick wanted to determine the length of one blade of the windmill pictured below. He stood at a point on the ground 440 feet from the windmill's base. Using surveyor's tools, Nick measured the angle between the ground and the highest point reached by the top blade and found it was 38.8° . He also measured the angle between the ground and the lowest point of the top blade, and found it was 30° .



Determine and state a blade's length, x , to the nearest foot.

$$\frac{440}{\sqrt{3}} = \frac{y\sqrt{3}}{\sqrt{3}}$$



$$\frac{440}{\sqrt{3}} \left(\frac{\sqrt{3}}{\sqrt{3}} \right) = y \left(\frac{\sqrt{3}}{\sqrt{3}} \right)$$

$$\frac{440\sqrt{3}}{3} = y$$


$$254.0341 = y$$

$$y = 254$$

Score 2: The student correctly found the height to the bottom of the top blade.

Question 34

34 Nick wanted to determine the length of one blade of the windmill pictured below. He stood at a point on the ground 440 feet from the windmill's base. Using surveyor's tools, Nick measured the angle between the ground and the highest point reached by the top blade and found it was 38.8° . He also measured the angle between the ground and the lowest point of the top blade, and found it was 30° .



Determine and state a blade's length, x , to the *nearest foot*.

$$\tan 30^\circ = \frac{x}{440}$$


$$x = -2818.3457 \text{ ft}$$

???

Score 1: The student wrote a correct trigonometric equation.

Question 34

34 Nick wanted to determine the length of one blade of the windmill pictured below. He stood at a point on the ground 440 feet from the windmill's base. Using surveyor's tools, Nick measured the angle between the ground and the highest point reached by the top blade and found it was 38.8° . He also measured the angle between the ground and the lowest point of the top blade, and found it was 30° .



Determine and state a blade's length, x , to the *nearest foot*.

$$\tan(8.8) = \frac{x}{440} =$$

$$x = (\tan(8.8))440$$


$$x = 68.115$$

the blade is
70 ft long

Score 0: The student gave a completely incorrect response.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{MEA}$, and $\overline{HA} \perp \overline{AT}$




Prove: $TA \cdot HA = HE \cdot TH$

STATEMENTS	REASONS
1. Quad $MATH$, $\overline{HE} \perp \overline{MEA}$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HA} \perp \overline{AT}$	1. Given
2. $\angle HEA$, $\angle TAH$ are rt \angle s	2. \perp lines form right \angle s.
3. $\angle HEA \cong \angle TAH$	3. All right \angle s are \cong .
4. $MATH$ is a \square	4. If a quadrilateral has 2 pairs of \cong opp sides, the quad is a \square .
5. $\overline{MA} \parallel \overline{TH}$	5. Opposite sides of a \square are \parallel .
6. $\angle THA \cong \angle EAH$	6. Alt int \angle s of \parallel lines and a transversal are \cong .
7. $\triangle HEA \sim \triangle TAH$	7. AA
8. $\frac{HA}{TH} = \frac{HE}{TA}$	8. Corresponding sides of similar \triangle s are in proportion.
9. $TA \cdot HA = HE \cdot TH$	9. In a proportion, the product of the means equals the product of the extremes.

Score 6: The student gave a complete and correct response.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{ME}$, and $\overline{HA} \perp \overline{AT}$



Prove: $TA \cdot HA = HE \cdot TH$

1. Quad $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$,
 $\overline{HE} \perp \overline{ME}$, $\overline{HA} \perp \overline{AT}$

2. $\overline{HA} \cong \overline{HA}$

3. $\triangle HAM \cong \triangle AHT$

4. $\angle THA \cong \angle MAH$

5. $\angle HEA$ & $\angle TAH$ are right \angle 's

6. $\angle HEA \cong \angle TAH$

7. $\triangle HEA \sim \triangle TAH$

$$8. \frac{HE}{TA} = \frac{HA}{TH}$$

$$9. TA \cdot HA = HE \cdot TH$$

1. Given

2. Reflexive

3. SSS \cong SSS

4. CPCTC

5. Perpendicular lines form right \angle 's

6. All right \angle 's are \cong .

7. AA \cong AA


8. Corresponding sides of similar triangles are in proportion.

9. The product of the means equals the product of the extremes.

Score 6: The student gave a complete and correct response.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{MEA}$, and $\overline{HA} \perp \overline{AT}$




Prove: $TA \cdot HA = HE \cdot TH$

Since $\overline{HE} \perp \overline{MEA}$ + $\overline{HA} \perp \overline{AT}$, $\angle HEA$ and $\angle TAH$ are right angles. $\angle HEA$ + $\angle TAH$ are \cong because all right angles are \cong . The opposite sides of quad. $MATH$ are \cong therefore $MATH$ is a \square so by definition $\overline{HT} \parallel \overline{MA}$. These parallel lines are cut by a transversal, \overline{HA} , which forms \cong alternate interior angles $\angle THA \cong \angle HAE$. $\triangle THA \cong \triangle HAE$ by $AA \cong AA$. In similar \triangle 's the corresponding sides are in proportion so $\frac{TA}{HE} = \frac{TH}{HA}$. In a proportion the cross products are equal therefore $TA \cdot HA = HE \cdot TH$.

Score 6: The student gave a complete and correct response.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{ME}$, and $\overline{HA} \perp \overline{AT}$




Prove: $TA \cdot HA = HE \cdot TH$

1) quad $MATH$, $\overline{MA} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{ME}$, $\overline{HA} \perp \overline{AT}$	2) Given	9) $TH \cdot EH = TA \cdot HA$ 9) In a proportion product of the means = the product of the extremes
2) quad $MATH$ is a \square	2) it has 2 sets of \cong oppo. sides	
3) $\overline{HT} \parallel \overline{MA}$	3) \square have 2 sets of \parallel sides	
4) $\angle THA \cong \angle HAE$	4) alt. int. \angle s are \cong if 2 sides are \parallel	
5) $\angle HEA$ and $\angle TAH = 90^\circ$	5) \perp lines form $90^\circ \angle$ s	
6) $\angle HEA \cong \angle TAH$	6) $90^\circ \angle$ s are all \cong	
7) $\triangle HEA \sim \triangle TAH$	7) $AA \sim$	
8) $\frac{TH}{TA} = \frac{HA}{HE}$	8) $\sim \triangle$ are proportional	

Score 5: The student had an incomplete reason in step 8.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{MEA}$, and $\overline{HA} \perp \overline{AT}$




Prove: $TA \cdot HA = HE \cdot TH$

Statements	Reasons
① $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{MEA}$, $\overline{HA} \perp \overline{AT}$	① Given
② $\angle 1 \cong \angle 2$	② All right angles are congruent.
③ $HMAT$ is a parallelogram	③ A parallelogram is a quad with both pairs opposite sides \cong
④ $\angle 3 \cong \angle 4$	④ Alternate interior angles formed by parallel lines \cong
⑤ $\triangle HEA \sim \triangle TAH$	⑤ AA \sim
⑥ $\frac{TA}{HE} = \frac{TH}{HA}$	⑥ Corresponding sides of similar triangles are in proportion
⑦ $TA \cdot HA = HE \cdot TH$	⑦ In a proportion, the product of the means is equal to the product of the extremes

Score 4: The student had two missing statements and reasons: stating angles 1 and 2 are right angles and stating $\overline{HT} \parallel \overline{MA}$.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{MEA}$, and $\overline{HA} \perp \overline{AT}$



Prove: $TA \cdot HA = HE \cdot TH$


We were given the opposite sides of quadrilateral $MATH$ are congruent, therefore $MATH$ is a Parallelogram. By definition the opposite sides \overline{HT} & \overline{AM} are parallel as well, with \overline{HA} as a transversal the alternate interior angles $\angle THA$ & $\angle HAE$ are congruent ($a \cong a$). Given $\overline{HE} \perp \overline{MEA}$ & $\overline{HA} \perp \overline{AT}$, right angles are formed, $\angle HEA \cong \angle TAH$ ($a \cong a$). By the reflexive property, $\overline{HA} \cong \overline{HA}$ ($s \cong s$). So by $AAS \cong AAS$ $\triangle HEA \cong \triangle TAH$, therefore $\triangle HEA \sim \triangle TAH$.

Since the corresponding sides of similar triangles are proportional the proportion $\frac{TA}{HE} = \frac{TH}{HA}$ can be derived, therefore $(TA)(HA) = (HE)(TH)$.

Score 3: The student incorrectly proved $\triangle HEA$ and $\triangle TAH$ congruent ($AAS \cong ASA$) and had a missing reason for $(TA)(HA) = (HE)(TH)$.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{ME}$, and $\overline{HA} \perp \overline{AT}$




Prove: $TA \cdot HA = HE \cdot TH$

Statements	Reasons
1.) $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{ME}$ and $\overline{HA} \perp \overline{AT}$	1.) Given
2.) $\angle 1$ & $\angle 2$ are rt \angle 's	2.) \perp lines form rt \angle 's
3.) $\angle 1 \cong \angle 2$	3.) all rt \angle 's are \cong
4.) $\overline{HA} \cong \overline{HA}$	4.) reflexive property

Score 2: The student correctly proved $\angle HEA \cong \angle TAH$, but step 4 is not relevant in proving $\triangle HEA \sim \triangle TAH$.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{ME}$, and $\overline{HA} \perp \overline{AT}$




Prove: $TA \cdot HA = HE \cdot TH$

S	R
① $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{ME}$, $\overline{HA} \perp \overline{AT}$	① Given
② $\angle HEA$, $\angle HAT$, $\angle MEH$ are 90°	② Perpendicular \rightarrow form 90°
③ $\angle HEA \cong \angle HAT \cong \angle MEH$	③ are congruent
④ $\overline{HA} \cong \overline{HA}$ ④.5 shape HAM is a parallelogram	④ reflexive property ④.5 definition of a parallelogram
⑤ $\angle THA \cong \angle EAH$	⑤ If 2 lines are cut by a transversal then all interior angles are \cong
⑥ $\triangle HMA \cong \triangle TAH$	⑥ SAS congruence
⑦ $TA \cdot HA = HE \cdot TH$	⑦ CPCTC

Score 2: The student wrote some correct relevant statements and reasons (steps 2 and 4).

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{ME}$, and $\overline{HA} \perp \overline{AT}$



Prove: $TA \cdot HA = HE \cdot TH$

1. $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$
 $\overline{HE} \perp \overline{ME}$, $\overline{HA} \perp \overline{AT}$

2. $\angle HEA$ + $\angle HAT$ are right \angle s

3. $\angle M \cong \angle T$

4. $\triangle HEA \cong \triangle HAT$

5. $\frac{TA}{HE} = \frac{TH}{HA}$

6. $TA \cdot HA = HE \cdot TH$

1. Given

2. Perpendicular lines form right \angle s

3. opposite angles \cong

4. AAS


5. CPCTC

6. Cross multiply

Score 1: The student had only one correct relevant statement and reason in step 2.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{MEA}$, and $\overline{HA} \perp \overline{AT}$




Prove: $TA \cdot HA = HE \cdot TH$

$\begin{array}{l} 1) \overline{HT} \cong \overline{AM}; \overline{HM} \cong \overline{AT} \\ 3) \triangle MHA \cong \triangle TAH \end{array}$	$\begin{array}{l} 2) \text{ Given} \\ 4) \text{ SSS} \end{array}$
--	---

Score 0: The student did not show enough correct relevant work to receive any credit.

Question 35

35 Given: Quadrilateral $MATH$, $\overline{HM} \cong \overline{AT}$, $\overline{HT} \cong \overline{AM}$, $\overline{HE} \perp \overline{MEA}$, and $\overline{HA} \perp \overline{AT}$



Prove: $TA \cdot HA = HE \cdot TH$

- | | |
|---|--|
| 1.) $\overline{HM} \cong \overline{AT}$ | 1.) Given |
| 2.) $\overline{HT} \cong \overline{AM}$ | 2.) Given |
| 3.) $\overline{HE} \perp \overline{MEA}$ | 3.) Given |
| 4.) $\overline{HA} \perp \overline{AT}$ | 4.) Given |
| 5.) $\sphericalangle 1, \sphericalangle 2,$ and $\sphericalangle 3$
are right \sphericalangle 's | 5.) def. right \sphericalangle 's |
| 6.) $\sphericalangle 1 \cong \sphericalangle 2$ | 6.) All right \sphericalangle 's are \cong |
| 7.) T is a reflexive \sphericalangle | 7.) Reflexive Property |
| 8.) $\triangle THA \sim \triangle HEA$ | 8.) AA \sim |
| 9.) $TA \cdot HA = HE \cdot TH$ | 9.) CPCTC |

Score 0: The student did not show enough correct relevant work to receive any credit.